

Brown dwarfs and very low mass stars in the Praesepe open cluster: a dynamically unevolved mass function?*

S. Boudreault¹, C. A. L. Bailer-Jones¹, B. Goldman¹, T. Henning¹ and J. A. Caballero²

¹ Max-Planck-Institut für Astronomie, Königstuhl 17, D-69117 Heidelberg, Germany
e-mail: boudrea, calj, goldman, henning@mpia.de

² Departamento de Astrofísica, Facultad de Física, Universidad Complutense de Madrid, E-28040 Madrid, Spain

Received 30 July, 2009; accepted 23 October, 2009

ABSTRACT

Context. Determination of the mass functions of open clusters of different ages allows us to infer the efficiency with which brown dwarfs are evaporated from clusters to populate the field.

Aims. In this paper we present the results of a photometric survey to identify low mass and brown dwarf members of the old open cluster Praesepe (age 590^{+150}_{-120} Myr, distance $190^{+6.0}_{-5.8}$ pc) from which we estimate its mass function and compare this with that of other clusters.

Methods. We performed an optical (I_c -band) and near-infrared (J and K_s -band) photometric survey of Praesepe covering 3.1 deg^2 . With 5σ detection limits of $I_c = 23.4$ and $J = 20.0$, our survey is predicted to be sensitive to objects with masses from 0.6 to $0.05 M_\odot$.

Results. We photometrically identify 123 cluster member candidates based on dust-free atmospheric models and 27 candidates based on dusty atmospheric models. The mass function rises from $0.6 M_\odot$ down to $0.1 M_\odot$ (a power law fit of the mass function gives $\alpha=1.8\pm0.1$; $\xi(M) \propto M^{-\alpha}$), and then turns over at $\sim 0.1 M_\odot$. This rise agrees with the mass function inferred by previous studies, including a survey based on proper motion and photometry. In contrast, the mass function differs significantly from that measured for the Hyades, an open cluster with a similar age ($\tau \sim 600$ Myr). Possible reasons are that the clusters did not have the same initial mass function, or that dynamical evolution (e.g. evaporation of low mass members) has proceeded differently in the two clusters. Although different binary fractions could cause the observed (i.e. system) mass functions to differ, there is no evidence for differing binary fractions from measurements published in the literature. Of our cluster candidates, six have masses predicted to be equal to or below the stellar/substellar boundary at $0.072 M_\odot$.

Key words. stars: low-mass, brown dwarfs – stars: luminosity function, mass function – stars: formation – Galaxy: open clusters and associations: individual: Praesepe

1. Introduction

Several publications in the past decade have been concerned with the mass function (MF) of low mass stellar and substellar populations in open clusters, including σ Orionis (Béjar et al. 2002, Caballero et al. 2007), the Orion Nebula Cluster (Hillenbrand & Carpenter 2000, Slesnick et al. 2004), IC 2391 (Barrado y Navascués et al. 2004, Boudreault & Bailer-Jones 2009), the Pleiades (Moraux et al. 2003, Lodieu et al. 2007), and the Hyades (Reid & Hawley 1999, Bouvier et al. 2008), to name just a few. Studies of relatively old open clusters (age ≥ 100 Myr) are important for the following two reasons in particular. First, they permit a study of the intrinsic evolution of brown dwarfs (BDs), e.g. their luminosity and effective temperature, which constrains structural and atmospheric models. Second, together with younger clusters we can investigate how BD populations as a whole evolve and thus probe the efficiency with which BDs evaporate from clusters to populate the Galactic field. Numerical simulations of cluster evolution have demonstrated that the MFs can evolve through dynamical interaction (de la Fuente Marcos & de la Fuente Marcos 2000; Adams et al. 2002b). These interactions result in a decrease of the open cluster BD (and low-mass star) population. This has been observed

by Bouvier et al. (2008) from a comparison of the Pleiades (120 Myr) and Hyades (625 Myr) mass functions.

Many earlier studies of the substellar MF have focused on young open clusters with ages less than ~ 100 Myr, and in many cases much younger (< 10 Myr). This is partly because BDs are bright when they are young (lacking a significant nuclear energy source, they cool as they age), thus easing detection of the least massive objects. However, youth presents difficulties. First, intra-cluster extinction plagues the determination of the intrinsic luminosity function from the measured photometry. Second, at these ages the BD models have large(r) uncertainties (Baraffe et al. 2002). Estimates of the substellar MF in very young clusters (age $\lesssim 1$ Myr) might be unreliable due to these modelling uncertainties (Chabrier et al. 2005). BDs in older clusters suffer less from these problems, but have the disadvantage that much deeper surveys are required to detect them.

The old open cluster Praesepe is an interesting target considering its age and distance. It is located at a distance of $190^{+6.0}_{-5.8}$ pc (based on parallax measurements from the new Hipparcos data reduction, van Leeuwen 2009) and has an age of 590^{+150}_{-120} Myr (by isochrone fitting in the Hertzsprung-Russell diagram; Fossati et al. 2008). The extinction towards this cluster is low, $E(B - V) = 0.027 \pm 0.004 \text{ mag}$ (Taylor 2006), while determinations of the metallicity of Praesepe yield some discrepancies: $[\text{Fe}/\text{H}] = 0.038 \pm 0.039$, Friel & Boesgaard (1992); $+0.13 \pm 0.10$, Boesgaard & Budge (1988); 0.11 ± 0.03 from spec-

* Based in part on observations carried out at ESO/La Silla, Chile under proposal number 078.A-9055(A).

troscopy and 0.20 ± 0.04 from photometry, An et al. (2007); $+0.27 \pm 0.10$, Pace et al. (2008). Hambly et al. (1995) presented a $\sim 19 \text{ deg}^2$ survey of the Praesepe cluster down to masses of $\sim 0.1 M_\odot$ and observed a rise of the MF at the lowest masses. They concluded that this implied a large population of BDs. A shallow survey complete to $I = 21.2 \text{ mag}$, $R = 22.2 \text{ mag}$ over 800 arcmin^2 uncovered one spectrally confirmed very low-mass star or BD (spectral type of M8.5V) with a model-dependent mass of $0.063\text{--}0.084 M_\odot$ (Magazzú et al. 1998). A survey over the central 1 deg^2 with 10σ limits of $R = 21.5$, $I = 20.0$ and $Z = 21.5 \text{ mag}$ revealed 19 BD candidates and the first MF determination of Praesepe down to the substellar limit, but without spectral confirmation (Pinfield et al. 1997). Subsequent infrared photometry of the sample reduced this number to nine candidates (Hodgkin et al. 1999). Adams et al. (2002a) presented a 100 deg^2 study of Praesepe using 2MASS (Two-Micron All Sky Survey) data and Palomar survey photographic plates, from which they derived proper motions. They determined the radial profile of this cluster but their MF does not reach the substellar regime. A more recent proper motion survey of Praesepe covers a much larger area (300 deg^2 ; Kraus & Hillenbrand 2007), but does not reach the BD regime either (the limit is $\sim 0.12 M_\odot$). Finally, the most recent substellar MF determination of Praesepe was published by González-García et al. (2006) and extends to a 5σ detection limit of $i = 24.5 \text{ mag}$ corresponding to $0.050\text{--}0.055 M_\odot$. They identified one new substellar candidate, but their survey covers only 1177 arcmin^2 .

In this paper, we present the results of a program to study, in detail, the MF of Praesepe down to the substellar regime. Our photometric survey is, as with González-García et al. (2006), the deepest so far in optical and near-infrared (NIR) bands, with 5σ detection limits of $I_c = 23.4$ and $J = 20.0$ (corresponding to a mass limit of about $0.05 M_\odot$), but covers more than nine times the area. Our paper is structured as follows. We first present the data set, reduction procedure and calibration in section 2. We then discuss our candidate selection procedure in section 3 and the survey results in section 4 before discussing the derived MF in section 5. We conclude in section 6.

2. Observations, data reduction, calibration, and estimation of masses and effective temperatures

2.1. Observations

Our survey consists of 47 Omega 2000 (O2k) fields each of size $15.4 \times 15.4 \text{ arcmin}^2$ observed in J and K_s , plus the same region observed in nine I_c Wide Field Imager (WFI) fields each of size $34 \times 33 \text{ arcmin}^2$. This gives a total coverage of 3.1 deg^2 observed in all three bands, centred on $\text{RA}(J2000) = 08^h 40^m 04^s$ and $\text{DEC}(J2000) = +19^\circ 40' 00''$.

The near-infrared (NIR) observations were made on the 3.5m telescope at Calar Alto, Spain (with observation runs of several nights from February 2005 to January 2007). O2k (Bailer-Jones et al. 2000; Baumeister et al. 2003) comprises a HAWAII-2 detector with $2k \times 2k$ pixels over a field of view of $15.4 \times 15.4 \text{ arcmin}$ delivering a pixel scale of 0.45 arcsec per pixel. The optical observations were carried out with the Wide Field Imager (WFI) on the MPG/ESO 2.2m telescope at La Silla, Chile (Baade et al. 1999) during 17–22 March 2007. The WFI is a mosaic camera of 4×2 CCDs, each with $2k \times 4k$ pixels, covering a total field of view of $34 \times 33 \text{ arcmin}^2$ at 0.238 arcsec per pixel. All fields were observed in the broad band filter I_c . A detailed list of the fields observed with pointing, filter, exposure time and 5σ detection limits is given in Table 1 for the NIR data and in

Table 1. Description of observations with the O2k infrared camera.

Field	α ($^h m s$)	δ ($^\circ ' ''$)	$t_{\text{exp}}(J)$ [min]	$t_{\text{exp}}(K_s)$ [min]	$J(5\sigma)$ [mag]	$K_s(5\sigma)$ [mag]
001	08 40 04	+19 40 00	60	40	19.9	18.6
A01	08 41 04	+19 54 00	60	130	20.6	19.4
A02	08 40 04	+19 54 00	40	40	19.5	18.5
A03	08 39 04	+19 54 00	40	40	19.7	18.7
A04	08 39 04	+19 40 00	20	40	19.0	18.6
A05	08 39 04	+19 26 00	20	40	20.3	18.8
A06	08 40 04	+19 26 00	20	40	19.9	18.6
A07	08 41 04	+19 26 00	40	60	20.3	19.0
A08	08 41 04	+19 40 00	20	40	18.0	17.6
B01	08 42 04	+20 08 00	40	40	19.7	18.4
B02	08 41 04	+20 08 00	40	40	18.8	18.6
B03	08 40 04	+20 08 00	20	40	20.0	18.5
B04	08 39 04	+20 08 00	20	40	19.7	18.6
B05	08 38 04	+20 08 00	20	40	20.3	18.7
B06	08 38 04	+19 54 00	20	40	20.0	18.5
B07	08 38 04	+19 40 00	20	40	19.5	18.5
B08	08 38 04	+19 26 00	20	40	17.4	17.8
B09	08 38 04	+19 12 00	20	40	18.0	18.8
B10	08 39 04	+19 12 00	20	40	20.1	18.8
B13	08 42 04	+19 12 00	20	40	20.1	18.8
B14	08 42 04	+19 26 00	20	40	20.1	18.7
B15	08 42 04	+19 40 00	20	40	19.3	18.3
B16	08 42 04	+19 54 00	20	40	18.4	18.1
C01	08 43 04	+20 22 00	20	40	20.4	18.8
C02	08 42 04	+20 22 00	20	40	20.1	18.7
C04	08 40 04	+20 22 00	20	40	20.0	18.8
C05	08 39 04	+20 22 00	20	40	19.1	18.2
C06	08 38 04	+20 22 00	20	40	20.1	18.7
C07	08 37 04	+20 22 00	20	40	19.8	18.2
C08	08 37 04	+20 08 00	20	40	19.9	18.7
C09	08 37 04	+19 54 00	20	40	20.1	18.0
C10	08 37 04	+19 40 00	20	40	20.3	19.3
C11	08 37 04	+19 26 00	20	40	19.6	18.6
C12	08 37 04	+19 12 00	20	40	20.3	18.9
C13	08 37 04	+18 58 00	20	40	20.4	18.7
C14	08 38 04	+18 58 00	20	40	19.9	18.4
C15	08 39 04	+18 58 00	20	40	20.4	18.6
C16	08 40 04	+18 58 00	20	40	19.4	18.4
C17	08 41 04	+18 58 00	20	40	19.3	18.3
C18	08 42 04	+18 58 00	20	40	20.6	18.3
C19	08 43 04	+18 58 00	20	40	20.3	18.4
C20	08 43 04	+19 12 00	20	40	19.6	18.6
C21	08 43 04	+19 26 00	20	40	20.3	18.8
C22	08 43 04	+19 40 00	20	40	20.3	18.7
C23	08 43 04	+19 54 00	20	40	19.8	18.7
C24	08 43 04	+20 08 00	20	40	19.1	18.4

Table 2 for the optical data. The passband functions for the filters, multiplied with the quantum efficiency of the detectors, are shown in Figure 1.


2.2. Data reduction and astrometry

The standard data reduction steps (overscan subtraction, trimming and flat-fielding for the WFI data; dark subtraction and flat-fielding for O2k data) were performed on a nightly basis, using the *ccdred* package under IRAF¹. For both WFI and O2k data we used *superflats* (obtained by combining science image

¹ The O2k camera suffers from a stray light problem. It appears on every image taken with the camera, forming a ring pattern centred in the middle of the detector (Nicol 2009). The stray light is removed via our global illumination correction and sky subtraction.

Table 2. Description of observations with WFI optical camera.

Field	α (^h ^m ^s)	δ ([°] ['] ^{''})	$t_{\text{exp}}(I_c)$ [min]	$I_c(5\sigma)$ [mag]
1	08 40 04	+19 40 00	24	22.6
2	08 42 24	+20 15 00	30	23.3
3	08 40 04	+20 15 00	36	23.2
4	08 37 44	+20 15 00	24	23.4
5	08 37 44	+19 40 00	36	23.6
6	08 37 44	+19 05 00	36	23.1
7	08 40 04	+19 05 00	36	23.5
8	08 42 24	+19 05 00	36	22.8
9	08 42 24	+19 40 00	36	22.8

**Fig. 1.** Transmission curve of the filters used in our survey compared to the synthetic spectrum of a BD with $T_{\text{eff}} = 2300$ K, $\log g = 4.5$ and solar metallicity (NextGen model). The transmission curves include the quantum efficiency of the detectors.

frames for each night) for pixel-to-pixel variation correction and for correcting the global illumination. For our NIR data, the sky background was subtracted using the median-combined images for each filter and each field (on a nightly basis). For WFI data, we reduced each of the eight CCDs in the mosaic independently and in the final step scaled them to a common flux response level. We made an initial sky subtraction via a low-order fit to the optical data, and for the infrared data by subtracting a median combination of all (unregistered) images of the science frames. Fringes were visible for the I_c -band photometry. They were removed in the way described by Bailer-Jones & Mundt (2001)². Finally, the individual images of a given field were registered and median combined. We used the IRAF task *daofind* to automatically detect stellar objects in an image by approximating the stellar point spread function with a Gaussian. We visually inspected the images in order to remove from our cluster candidate list any extended sources (i.e. galaxies) that were mistakenly identified as stars by *daofind* (see section 3.3). Sources were extracted and instrumental magnitudes assigned via aperture photometry with the IRAF task *wphot*. To this aperture photometry we have ap-

² A *fringe correction frame* was created, which is a median combination of all science frames in a same filter with the same exposure time. This *fringe correction frame* was scaled by a factor, determined manually for each science frame, and subtracted from the science image.

plied an aperture correction following the technique described in Howell (1989). An astrometric solution was obtained using the IRAF package *imcoords* and the tasks *ccxymatch*, *ccmap* and *cctran*. For each WFI field, this solution was computed for the I_c -band image (and for each O2k field using the J -band image) using the 2MASS catalogue as a reference. The root mean square accuracy of our astrometric solution is 0.15–0.20 arcsec for both WFI and O2k data. For WFI data, the astrometry was performed on a CCD-by-CCD basis.

2.3. Photometric calibration

To correct for Earth-atmospheric absorption on the photometry, we calibrated the infrared data using the J and K_s -band magnitudes of 2MASS objects which were observed in our science fields. By determining a constant offset between the magnitude of 2MASS and our instrumental magnitude, we obtained the zero point offset. Since this zero point offset was obtained with objects in the same field of view in each science frame, and since we found the difference between the 2MASS and O2k passbands to be insignificant, we did not need to perform an airmass or colour correction when reducing our NIR photometry. (That is, the determined coefficients were statistically consistent with zero.)

We followed a similar approach for our I_c -band photometry, but using observations in our fields for which r and i -band magnitudes are available in the Sloan Digital Sky Survey (SDSS) catalogue. We first transformed the i -band magnitudes of SDSS to I_c -band magnitudes using the transformation equation of Jordi et al. (2006)

$$I_c = i - 0.381 - 0.254 \times (r - i) \quad (1)$$

We then determined the zero point offset between this I_c magnitude and our instrumental I_c magnitude, again using the SDSS stars. A further colour correction was not necessary, and as this calibration is applied on a field-by-field basis (as with the NIR data), an airmass correction was likewise unnecessary.

2.4. Mass and effective temperature estimates based on photometry

After we identify candidates (section 3) we will use the multi-band photometry to derive their masses and effective temperatures, T_{eff} . We use the evolutionary tracks from Baraffe et al. (1998) and atmosphere models from Hauschildt et al. (1999a) (assuming a dust-free atmosphere; the NextGen model) to compute an isochrone for Praesepe for an age of 590 Myr, a distance of 190 pc, a solar metallicity and assuming zero extinction. These models and assumptions provide us with a prediction of f_λ , the spectral energy distribution received at the Earth (in $\text{erg cm}^{-2} \text{s}^{-1} \text{\AA}^{-1}$) from the source. We need to convert these spectral energy distributions into magnitudes in the filters we used. Denoting as $S_A(\lambda)$ the (known) total transmission function of filter A (including the CCD quantum efficiency and assuming telescope and instrumental throughput are flat), then the flux measured in the filter is

$$f_A = \frac{\int_0^\infty f_\lambda S_A(\lambda) d\lambda}{\int_0^\infty S_A(\lambda) d\lambda}, \quad (2)$$

The corresponding magnitude m_A in the Johnson photometric system is given by

$$m_A = -2.5 \log f_A + c_A, \quad (3)$$

where c_A is a constant (zero point) that remains to be determined in order to put the model-predicted magnitude onto the Johnson system. We derived this constant for each of the bands I_c , J and K_s in the standard way, namely by requiring that the spectrum of Vega produce a magnitude m_A of 0.03 in all bands. Using the Vega spectrum from Colina et al. (1992) we derive values of $c_{I_c} = -22.6011$, $c_J = -23.6865$ and $c_{K_s} = -25.9076$ mag. Applying the two equations above to a whole set of model spectra produces a theoretical isochrone in colour–magnitude space. Note that this procedure only provides us with the “true” magnitudes of the model spectra, not their instrumental ones. The photometric calibration procedure applied to the data converts the measured, instrumental magnitudes to the “true” magnitude plane where we then compare them with the isochrone.

Assuming that all our photometric candidates belong to Praesepe, we derive masses and effective temperatures from these isochrones using our three filter measurements in the following way. We first normalize the energy distribution of each object to the energy distribution of the model using the J filter. We then estimate the mass and effective temperature via a least squares fit of the measured spectral energy “distribution” (actually just two points) to the model spectral energy distribution from the isochrone. This involves estimating one parameter from two measurements, because mass and T_{eff} are not independent.

The above assumption of a dust-free atmosphere is valid for $T_{\text{eff}} \geq 3000$ K, but objects with $3000 \text{ K} \geq T_{\text{eff}} \geq 1800$ K are expected to have dust in equilibrium with the gas phase (Allard et al. 2001). We therefore perform a second selection of candidates (and determination of mass and T_{eff}) based on isochrones predicted in the same way, but based on evolutionary tracks of Chabrier et al. (2000) and the AMES-dusty model of Allard et al. (2001). This gives us a second *dusty model* list of candidates. A priori some observed stars could appear in both lists (and in fact two do), but in our later discussions of the mass function we do not mix stars from the two lists but rather make separate determinations of the mass function.

There are various sources of error in the estimation of mass and T_{eff} . These are the photon noise, the photometric calibration, the least squares fitting (imperfect model) and the uncertainties in the age of and distance to Praesepe. The uncertainties in the age and distance are the most significant errors and given rise to uncertainties of $0.060 \pm 0.010 M_\odot$ and 1990 ± 260 K for a substellar object, $0.072 \pm 0.008 M_\odot$ and 2293 ± 201 K for an object at the hydrogen burning limit and $1.000 \pm 0.017 M_\odot$ and 5300 ± 50 K for a solar-type star.

3. Candidate selection procedure

The candidate selection procedure for BDs and very low-mass stars is as follows (and explained in more detail in the remainder of this section). Candidates were first selected based on colour-magnitude diagrams (CMDs) and this further refined using colour-colour diagrams. In the third and final selection, we used the known distance to Praesepe to reject objects based on a discrepancy between the observed magnitude in J and the magnitude in this band computed with the isochrones and our estimation of T_{eff} . To be considered as a cluster member, an object has to satisfy all three of these criteria. We make two independent selections: one using dust-free and one using dusty atmospheric models.




Fig. 2. Colour–magnitude diagram showing an example of the first selection step using the I_c and J bands. The solid lines show the isochrone computed from an evolutionary model with a dust-free atmosphere (NextGen model) and the dashed lines show our selection band around this. The numbers indicate the masses (in $M_\odot$) of objects on the isochrone for various I_c magnitudes. Overplotted are measurements from our survey of candidate cluster members reported in Pinfield et al. (1997) (*stars*), Adams et al. (2002a) (*triangles*) [where we include objects which have a probability of being a real member higher than 10%], González-García et al. (2006) (*squares*) and Kraus & Hillenbrand (2007) (*circles*).

3.1. First candidate selection step: colour–magnitude diagrams

Candidates were first selected from our CMDs by retaining only objects which are no more than 0.14 mag redder or bluer than the isochrone in all CMDs. This number accommodates errors in the magnitudes, uncertainties in the model isochrones plus uncertainties in the cluster age and distance estimates. We additionally include objects brighter than the isochrones by 0.753 mag in order to include unresolved binaries. In Figure 2 and 3 we show two CMDs where candidates were selected based on I_c vs. $I_c - J$ and K_s vs. $I_c - K_s$. These figures also show low-mass cluster member candidates from previous studies which we detected in our survey (Pinfield et al. 1997; Adams et al. 2002a; González-García et al. 2006; Kraus & Hillenbrand 2007). In Figure 3, we can observe three structures in this CMD. The two structures at $I_c - K_s \sim 1$ mag and $I_c - K_s \sim 2$ mag are predominantly stars (Galactic disk turn-off, and disk late-type and giant stars respectively) while the structure at $I_c - K_s \sim 3$ mag is mostly composed of galaxies. From a total of 23 891 objects detected above the 5σ detection limit in all filters, 800 are retained as candidate cluster members (96.7% are rejected). If we instead use dusty model isochrones, then out of the 23 891 objects, 357 are retained (98.5% are rejected) for our dusty model list.




Fig. 3. As Figure 2 but with the I_c and K_s bands.

3.2. Second candidate selection step: colour-colour diagram

The second stage of candidate selection involves retaining just those objects which lie within 0.24 mag of the isochrone in the (single) colour-colour diagram. This value reflects the photometric errors plus uncertainty in the age estimation of Praesepe. One such colour-colour diagram with the selection limits is shown in Figure 4. The two main sources of contamination beside field M dwarfs are background red giants and unresolved galaxies (Praesepe is at a Galactic latitude of $b=+32.5^\circ$). We show in Figure 5 the theoretical colours for red giants using the atmosphere models of Hauschildt et al. (1999b) and theoretical colours of six galaxies from Meisenheimer et al. (2009). We see that red giants could be a source of contamination in the mass range of $0.09\text{--}0.2\text{ M}_\odot$ and at $\sim 0.7\text{ M}_\odot$, while unresolved galaxies should not be a major source of contamination below $0.6\text{ M}_\odot$. In Figure 5 we see the same three structures as in Figure 3: from top to bottom galaxies, disk late-type and giant stars, and Galactic disk turn-off stars. Of the 800 objects selected in the first step, 291 are kept here (63.6% are rejected) assuming a dust-free atmosphere, and 110 out of 357 are kept (69.2% are rejected) when using the model for a dusty atmosphere.

3.3. Third candidate selection step: Rejection based on observed magnitude vs. predicted magnitude discrepancy

As indicated in section 2.4, our determination of T_{eff} is based on the spectral energy distribution of each object and is independent of the assumed distance. The membership status of an object can therefore be assessed by comparing its observed magnitude in a band with its magnitude predicted from its T_{eff} and Praesepe's isochrone (which assumes a distance). The premise is that the predicted magnitude of a background contaminant would be lower (brighter) than its observed magnitude and higher (fainter) for a foreground contaminant. In order to avoid removing unresolved binaries that are real members of the cluster, we keep all objects with a computed magnitude of up to




Fig. 4. Colour-colour diagram used in the second selection step. The solid line is the isochrone computed from an evolutionary model with a dust-free atmosphere (NextGen model, the masses in $\text{M}_\odot$ for each $I_c\text{--}J$ colours are shifted up clarity) and the dashed lines show our selection band. Overplotted are the cluster candidate members from Pinfield et al. (1997) (stars), Adams et al. (2002a) (triangles), González-García et al. (2006) (squares) and from Kraus & Hillenbrand (2007) (circles), which we detected in our survey.




Fig. 5. As Figure 4, but now showing the theoretical colours of six galaxies as thick dotted lines and the theoretical colours of red giants as thick solid lines. The six galaxies are two starbursts, one Sab, one Sbc, and two ellipticals of 5.5 and 15 Gyr, with redshifts from $z=0$ to $z=2$ in steps of 0.25 (evolution not considered). We assume that all red giants have a mass of $5\text{ M}_\odot$, $0.5 < \log g < 2.5$ and $2000\text{ K} < T_{\text{eff}} < 6000\text{ K}$.




Fig. 6. Difference between the observed J magnitude and the model J magnitude computed from the derived mass and T_{eff} , as a function of T_{eff} . The two vertical lines are at the positions of L0 and M5 dwarfs (left to right), for orientation purposes. The dotted line (at -0.753 mag) represents the offset due to the possible presence of unresolved binaries, the dot-dash line represents the error on the magnitude determination, and the curved, long-dashed lines represent the uncertainties on the age and distance of Praesepe.

0.753 mag brighter than the observed magnitude. We also take into account photometric errors and uncertainties in the age and distance of Praesepe. This selection procedure is illustrated in Figure 6. From 291 objects selected through CMDs and colour-colour diagrams in the first two steps, 144 are kept (50.5% are rejected) when using the dust-free atmospheres/models, and 35 out of 110 are kept (68.2% are rejected) when using the dusty atmosphere/models.

After this step, we perform a visual inspection directly on the images to reject resolved galaxies and spurious detections. This inspection removes 21 and 8 objects from the dust-free and dusty selection respectively.

4. Results of the survey

We now present the selected candidates, discuss contamination by cluster non-members and derive the magnitude and mass functions for Praesepe.

4.1. Selected photometric candidates

The final selection reveals 123 photometric candidates using an isochrone based on dust-free atmospheres, and 27 objects using an isochrone assuming dusty atmospheres³. This corresponds to ~ 40 and ~ 9 objects per deg² respectively. All our photometric candidates are presented in Table 3. Objects are given the notation PRAESEPE-YYY where YYY is a serial identification number (ID). Numbers above 900 indicate candidate members

³ Two objects in the *dust-free atmosphere selection* (PRAESEPE-089 and -093) were also identified in *dusty atmosphere selection* (PRAESEPE-915 and -917).

Table 4. Photometric candidates in our survey that are also photometric candidates in previous surveys.

ID	α (^h ^m ^s)	δ ([°] ' ")	Alternative name	Ref. ^a
005	08 41 08.5	+19 54 02.0	RIZpr18	[3]
010	08 39 06.9	+19 47 08.0	J0839069+194708	[1]
			J08390695+1947080	[2]
011	08 38 55.46	+19 50 33.3	J0838554+195033	[1]
			J08385547+1950334	[2]
012	08 38 54.19	+19 51 44.6	J08385420+1951446	[2]
			J0839030+192415	[1]
			J08390308+1924155	[2]
015	08 39 12.71	+19 30 16.8	J08391272+1930169	[2]
016	08 39 54.39	+19 27 37.1	J0839544+192737	[1]
			J08395441+1927372	[2]
017	08 39 47.82	+19 28 03.1	RIZpr11	[3]
029	08 42 50.50	+20 20 03.8	J0842505+202004	[1]
			J08425052+2020039	[2]
034	08 42 54.58	+20 03 36.3	RIZpr23	[3]
035	08 42 51.96	+20 05 19.4	J0842519+200519	[1]
038	08 43 10.76	+20 01 29.3	J0843107+200129	[1]
040	08 41 11.05	+20 22 38.4	J0841110+202238	[1]
042	08 40 10.59	+20 20 50.4	J0840106+202050	[1]
			J08401060+2020505	[2]
045	08 39 14.51	+20 01 19.1	J0839145+200119	[1]
046	08 39 22.43	+20 04 54.6	J0839224+200454	[1]
			J08392244+2004548	[2]
047	08 38 55.15	+20 13 08.8	J0838551+201308	[1]
			J08385517+2013089	[2]
054	08 40 53.96	+20 05 24.3	J08405397+2005243	[2]
062	08 36 39.46	+20 22 33.8	J08363947+2022339	[2]
064	08 36 44.99	+20 08 45.7	J08364501+2008459	[2]
066	08 37 11.41	+20 13 45.8	J08371143+2013459	[2]
068	08 38 08.0	+20 03 50.1	J08380800+2003505	[2]
070	08 38 12.44	+20 08 02.5	J08381244+2008026	[2]
073	08 38 21.85	+20 05 35.7	J08382186+2005356	[2]
075	08 38 39.27	+19 41 40.1	J08383929+1941401	[2]
081	08 37 24.48	+19 47 11.9	J08372449+1947120	[2]
082	08 37 02.1	+19 52 07.3	RIZpr2	[3]
101	08 41 20.32	+18 57 42.9	J08412034+1857430	[2]
103	08 42 19.21	+19 02 14.8	J08421923+1902148	[2]
108	08 43 09.0	+19 43 11.9	J08430905+1943119	[2]
109	08 43 01.2	+19 49 59.8	RIZpr24	[3]
117	08 42 11.47	+19 52 50.2	RIZpr21	[3]
110	08 43 01.9	+19 54 04.5	J08430186+1954046	[2]
112	08 42 52.26	+19 51 45.9	J08425228+1951460	[2]
116	08 42 15.48	+19 48 57.6	J08421550+1948576	[2]
122	08 43 08.4	+19 28 06.1	J08430839+1928061	[2]
123	08 43 12.63	+19 34 28.9	J08431265+1934290	[2]

^a Objects [1] are from Adams et al. (2002a), [2] are from Kraus & Hillenbrand (2007) and [3] are from Pinfield et al. (1997).

assuming a dusty atmosphere. Only the first 10 rows of the tables are shown, all other data are available online. We also note in Table 4 which objects are candidate cluster members also identified as such by Kraus & Hillenbrand (2007), Adams et al. (2002a) or Pinfield et al. (1997).

Some Praesepe members from previous studies are not detected in our work. This is the case for the objects from Pace et al. (2008) and Fossati et al. (2008), for example. Since those studies focused on bright objects, these stars saturate in our science images. (Pace et al. 2008 and Fossati et al. 2008 were concerned with chemical abundances of A-type and solar-type stars, respectively, while our saturation occurs at $\sim 0.7 M_{\odot}$.)

Table 3. All photometric cluster member candidates of our survey. Table 3 is published in its entirety in the electronic edition of *Astronomy & Astrophysics*. A fraction is shown here for guidance regarding its form and content.

ID	α (^h ^m ^s)	δ ([°] ['] ^{''})	I_c^a [mag]	J^a [mag]	K_s^a [mag]	M^a [$M_\odot$]	T_{eff}^a [K]	J_{model}^a [mag]
001	08 40 53.61	+19 40 58.6	19.19	16.81	15.61	0.089	2665	17.00
002	08 41 08.8	+19 43 27.5	16.14	14.95	13.97	0.219	3321	14.86
003	08 41 01.6	+19 52 02.5	16.67	15.35	14.38	0.162	3189	15.48
004	08 41 12.17	+19 52 48.6	18.43	16.39	15.46	0.099	2805	16.72
005	08 41 08.5	+19 54 02.0	19.02	16.58	15.39	0.088	2636	17.06
006	08 40 10.74	+19 40 49.8	16.97	15.47	14.36	0.132	3061	15.95
007	08 39 39.56	+19 47 54.3	17.95	16.10	15.07	0.104	2860	16.58
008	08 39 43.38	+19 48 45.7	16.89	15.56	14.68	0.161	3186	15.50
009	08 39 55.84	+19 53 14.3	20.29	17.50	16.54	0.081	2520	17.32
010	08 39 06.9	+19 47 08.0	16.51	15.14	14.20	0.155	3166	15.57

^a The 1σ uncertainty in the determination of magnitude, effective temperature and mass are the following : $\Delta\text{mag} = 0.002$ mag, $\Delta T_{\text{eff}} = 140$ K and $\Delta M = 0.1 M_\odot$ for stars ($M > 0.2 M_\odot$), $\Delta\text{mag} = 0.01$ mag, $\Delta T_{\text{eff}} = 230$ K and $\Delta M = 0.05 M_\odot$ for very low-mass stars ($0.072 M_\odot < M < 0.2 M_\odot$), $\Delta\text{mag} = 0.04$ mag, $\Delta T_{\text{eff}} = 420$ K and $\Delta M = 0.02 M_\odot$ for BDs ($M < 0.072 M_\odot$). The magnitude J_{model} is the predicted magnitude based on photometric determination of T_{eff} and mass.

Not all objects identified by other surveys as brown dwarfs or very low mass stellar member candidates – and detected in our survey – are members based on our criteria. The two objects from the work of González-García et al. (2006), who also used photometry in order to select candidate members, we detect above our 5σ limit (Prae J084039.3+192840 and Prae J084130.4+190449). Yet both objects are non-members based on our selection criteria, because they have $I_c - J$ colours bluer than our selection band. (Prae J084130.4+190449 is also too blue in $I_c - K_s$ for our selection band at $I_c - K_s = 3.0$ mag, whereas Prae J084039.3+192840 at $I_c - K_s = 4.0$ mag lies within it.) González-García et al. (2006) did not report any NIR photometry for these two objects. Although the non-membership of Prae J084039.3+192840 can be debated (high membership probability based on González-García et al. 2006), Prae J084130.4+190449 is most likely an unresolved galaxy (low membership probability; González-García et al. 2006).

Of the candidates from the photometric survey of Pinfield et al. (1997), seven fall within our survey and are detected, of which six are identified as candidates by our selection criteria. The non-selected object is RIZpr6 in Hodgkin et al. (1999). It is bluer than the isochrones in both CMDs in Figure 2 and 3. From its positions in the CMDs and in the colour-colour diagram in Figure 4, we suspect that this object is an unresolved galaxy.

11 of the the 14 objects from a survey based on proper motion and photometry by Adams et al. (2002a) are identified by our selection. The objects not recovered fail the observed magnitude vs. predicted magnitude test. On the other hand, 27 cluster candidates of Kraus & Hillenbrand (2007) out of 37 detected in our survey are selected. The 10 non-selected objects have membership probabilities from Kraus & Hillenbrand (2007) based on proper motion greater than 95%, and are brighter than the 10σ detection limit of the publicly available surveys used in their work. However, these objects failed our observed magnitude vs. predicted magnitude test and some are bluer than our isochrone of Praesepe in $J - K_s$. With $I_c - K_s$ colour of ~ 2 mag, we suggest that these objects are more likely to be disk late-type stars or giant stars.

The 5σ detection limits of our survey are $I_c=23.4$ mag, $J=20.0$ mag and $K_s=18.6$ mag (which correspond to $\sim 0.05 M_\odot$ using the dust-free isochrone). However, we cannot expect to detect *all* objects down to these magnitudes. We estimate the




Fig. 7. Estimation of the completeness limit for our survey using the J band. The solid line is the best linear fit before the turn off, the vertical dashed line is the 5σ detection limit and the vertical dotted line is the magnitude at which detector saturation occurs.

survey completeness by taking the ratio of the number of objects detected to the predicted number of detections, the latter made by assuming a uniform distribution of stars along the line of sight in our survey fields. (This comparison distribution is somewhat crude, but it gives an approximate value without making too many assumptions.) The predicted number of detections is obtained from the histogram of the number of detections as a function of magnitude (Figure 7) and by observing where the distribution drops off compared to a straight line extrapolation. Based on this, the completeness of the survey down to the 5σ detection limit is 90% in I_c , 88% in J and 87% in K_s . The overall detection completeness of our survey, from saturation to 5σ detection corresponding to $0.05 M_\odot$, is therefore $\sim 87\%$. In J band, we reach a completeness of 95% at $J=19.7$ mag, which corresponds to $\sim 0.055 M_\odot$.




Fig. 8. Finding charts of the six new BD candidates of Praesepe (J -band). We observed objects very close to PRAESEPE-099 and -909, although they do not influence the photometry. The panels are 50×50 arcsec with North up and East to the left.

4.2. Substellar candidates in Praesepe

Six objects in our survey are cluster candidates with theoretical masses equal to or below the stellar/substellar boundary at $0.072 M_{\odot}$. We present the finding charts of the six objects in Figure 8. In Table 5, we present their coordinates and physical parameters. These BD candidates have predicted masses between 0.064 and $0.072 M_{\odot}$. A spectroscopic follow up (on a 8 m class telescope or larger) will be needed in order to confirm or refute their membership and their substellar status.

4.3. Contamination by non-members

As mentioned in section 3.2, the two main sources of contamination are the background red giants, which are the dominant source at masses of 0.09 – $0.2 M_{\odot}$ and $\sim 0.7 M_{\odot}$, and unresolved galaxies, mostly affecting masses above $0.6 M_{\odot}$. Other possible contaminants are field M dwarfs and high redshift quasars (for instance at $z \sim 6$; Caballero et al. 2008). However, as such quasars have spectral energy distributions similar to mid-T dwarfs whereas our faintest candidates are early L dwarfs, and given that they are rare (3.3 quasars at $5.5 < z < 6.5$ in a 8 deg^2 survey, Stern et al. 2007), the MF should not be affected by quasar contamination.

Let us estimate the contamination by M dwarfs. First, we consider that close to the open cluster Praesepe, the space density of M dwarfs is *uniform*. We assume that their density (ρ) drops exponentially with vertical distance from the galactic disk (h) such that

$$\rho(h) = \rho_0 e^{-\frac{h}{h_0}}, \quad (4)$$

assuming a scale height of $h_0 = 500 \text{ pc}$. We use the local space density (ρ_0) for M dwarfs of $57 \cdot 10^{-3} \text{ pc}^{-3}$ (from the *Research Consortium on Nearby Stars*; Henry et al. 2006). Given the Galactic latitude of Praesepe of $b = +32.5 \text{ deg}$ and its distance of 190 pc , the density of M dwarfs near Praesepe should be $47 \cdot 10^{-3} \text{ pc}^{-3}$. If we define a volume corresponding to the area of our survey (3.1 deg^2 , 34 pc^2) and use the distance uncertainties to the cluster ($190^{+6.0}_{-5.8} \text{ pc}$) as its depth, we estimate that we have ~ 19 M dwarf contaminants near the cluster. From a total of 150 photometric candidates, we estimate a contamination of $\sim 13\%$ (or even less, as the cluster depth is presumably closer to




Fig. 9. J band luminosity function. The solid line histogram represents the luminosity function based on a selection using a dust-free atmosphere (NextGen model); the thick dotted histogram uses a dusty atmosphere (AMES-Dusty model). The stellar/substellar limit is at $J \sim 17.8 \text{ mag}$ for both models ($0.072 M_{\odot}$). For reference, the ordinate value of 1.11 at the histogram peak (magnitude bin $J = 15.25 \text{ mag}$) corresponds to 13 objects.

$\sqrt{34} = 5.8 \text{ pc}$ than to the 11.8 pc error span of the mean cluster distance). Therefore, we do not expect contamination by field M dwarfs to play a significant role in the determination of the MF.

4.4. Luminosity function and mass function

We present in Figure 9 the luminosity function of Praesepe using the J -band magnitude of the cluster candidates. No correction is made for binaries, so this is the system rather than single-star luminosity function.

The mass function (MF), $\xi(\log_{10} M)$, is generally defined as the number of stars per cubic parsec in the logarithmic mass interval $\log_{10} M$ to $\log_{10} M + d \log_{10} M$. Here, we do not compute the volume of Praesepe so instead we define the MF as the total number of objects in each $0.1 \log_{10} M$ bin per square degree. Since we do not make any corrections for binaries we compute here a *system* MF. Our inferred MF is shown in Fig. 10. The log-normal form for a MF is

$$\xi(\log_{10} M) = k \cdot \exp \left[-\frac{(\log_{10} M - \log_{10} M_0)^2}{2\sigma^2} \right], \quad (5)$$

where $k=0.086$, $m_0=0.22 M_{\odot}$ and $\sigma=0.57$ for the Galactic field system MF (Chabrier 2003). Fitting this function to both the *dust-free* and *dusty* MF data we obtain $k=5.9 \pm 3.1$, $m_0=0.15 \pm 0.05 M_{\odot}$ and $\sigma=0.51 \pm 0.12$. Figure 10 shows this result. If we instead fit a power law (Salpeter (1955))

$$\xi(\log_{10} M) = k \cdot M^{-\alpha}, \quad (6)$$

from the highest mass bin to the turn over at $0.1 M_{\odot}$, we obtain $\alpha=1.3 \pm 0.2$ (corresponding to $\xi(M) \propto M^{-2.3}$). If we exclude the two bins between 0.1 and $0.15 M_{\odot}$ (possible contamination by red giants) and the two highest bins (possible incompleteness), the fit gives $\alpha=0.8 \pm 0.1$.

Table 5. Same as Table 3, but only the BD candidates are given and we include the spectral type expected.

ID	α ($^h m s$)	δ ($^{\circ} ' ''$)	I_c [mag]	J [mag]	K_s [mag]	M [$M_{\odot}$]	T_{eff} [K]	J_{model} [mag]	SpT ^a
055	08 41 04.5	+20 14 58.0	21.61	18.29	17.12	0.068	2250	17.97	L0
096	08 41 13.48	+18 59 05.1	21.06	17.85	16.82	0.072	2335	17.75	M9
099	08 41 45.16	+19 18 07.7	21.30	17.98	17.01	0.068	2249	17.98	L0
909	08 39 29.94	+20 11 40.3	20.11	17.63	16.67	0.069	2259	17.95	L0
910	08 40 34.00	+20 14 56.2	20.08	17.60	16.65	0.069	2261	17.94	L0
915	08 38 51.77	+19 00 21.6	20.28	17.67	16.68	0.068	2238	18.01	L0


^a Spectral type expected based on T_{eff} and the temperature scale of Kraus & Hillenbrand (2007), with L1 set to 2100 K. The error on this estimation is one subclass.

Fig. 10. Mass function based on our survey photometry. Points with error bars represent the MF based on a selection and mass calibration assuming a dust-free atmosphere, whereas the open circles with error bars are the MF based on the dusty atmosphere model. We also overplot the log-normal and the power law MF fitted to our data (both solid line). Error bars are Poissonian arising from the number of objects observed in each bin. The vertical thin dotted lines are the mass limits at which detector saturation occurs in the I_c , J and K_s -bands (from left to right). The vertical thick dashed line is the mass at the 5σ detection limit (completeness of $\sim 87\%$). For reference, the ordinate value of 0.932 at the histogram peak (mass $\log_{10} M = -0.85$ [$0.142 M_{\odot}$]) corresponds to 27 objects. The two dusty data points have been shifted to the right by $\log_{10} M = 0.05$ for clarity.

5. Analysis and discussion of the stellar and substellar mass function of Praesepe

Our MF of Praesepe (Figure 10) shows a rise in the number of objects from $0.6 M_{\odot}$ down to $0.1 M_{\odot}$, and then a turn-over at $\sim 0.1 M_{\odot}$. This turn-over is not due to incompleteness since it occurs well above the 5σ detection limit corresponding to $0.05 M_{\odot}$. This behaviour is confirmed by the luminosity function in Figure 9 which shows a rise from $J=13$ to 16 mag (with candidates obtained using a dust-free atmosphere) and a drop at $J=17$ mag

(seen with both types of candidates). To help the analysis of these features in the mass function, we compare in Figure 11 the mass functions of Praesepe obtained from several studies plus the MF for the old open cluster Hyades (age of 625 Myr).

The rise in our MF of Praesepe is also present in the MFs obtained in the three previous studies of Baker & Jameson (2009), Kraus & Hillenbrand (2007) and Hambly et al. (1995). On the other hand, we do not see this rise in the MF of Adams et al. (2002a). However, their MF is based on objects with a membership probability higher than only 1% and within a radius of 3.8 deg. Due to use of such a low probability threshold for selection, we expect that most of the objects used in the MF determination are simply field stars (which is their own conclusion in section 5.4; Adams et al. 2002a), so further comparison is not warranted. As for the MFs of González-García et al. (2006) and Pinfield et al. (1997), since the highest mass bins are ~ 0.11 and $\sim 0.15 M_{\odot}$ (respectively), the rise observed from $0.6 M_{\odot}$ to $0.1 M_{\odot}$ cannot be discussed.

Only four MFs, in addition to our work, reach masses below $0.1 M_{\odot}$: Baker & Jameson (2009), González-García et al. (2006), Pinfield et al. (1997) and Hambly et al. (1995). While the MFs of Baker & Jameson (2009) and Hambly et al. (1995) show a turn-over at $0.1 M_{\odot}$, the one obtained by Pinfield et al. (1997) does not. On the contrary, it presents a sudden rise at the stellar/substellar limit (with a ratio of ~ 5 in the number of objects at the mass bin at 0.07 to the number in the bin at $0.11 M_{\odot}$). They used *RIZ* photometry for their survey, but not all objects were observed in all bands, resulting in just one colour available for membership determination in some cases (Pinfield et al. 1997). From an analysis of MFs of other clusters and using a multi-band photometric survey, Boudreault & Bailer-Jones (2009) have shown that use of a narrow spectral coverage with few filters can lead to high contamination by field M dwarfs, and thus an apparent rise in the MF in the low mass regime. We suggest that this is the reason for the apparent rise at the low-mass end of the MF in Pinfield et al. (1997) (who also noted that only one colour is available for many objects in their two lowest bins). As for the MF of González-García et al. (2006), as they only have three points we cannot comment on any possible trend.

Although there are some discrepancies between the different MFs of Praesepe from previous works and our MF, none agrees with the MF of the Hyades (~ 625 Myr) obtained by Bouvier et al. (2008)⁴, in which the MF is observed to turn-over and decrease already at $0.35 M_{\odot}$. This is surprising, since Praesepe and the Hyades share a com-

⁴ Like the MF of Praesepe we present, the MF of the Hyades presented by Bouvier et al. (2008) is a system MF (no correction for binaries).




Fig. 11. MF of Praesepe from our present work (*open dots* assuming a dusty atmosphere and *filled dots* assuming a dust-free atmosphere), from previous work (*open triangles* for survey using proper motion and *filled triangles* for survey using photometry only), as well as the MF from the Hyades from Bouvier et al. (2008) (*open squares*). We also show the galactic field star MF from Chabrier (2003) as a thin dashed line and the substellar limit as a thick dashed line. We have normalized all the MFs to the log-normal fit of Chabrier et al. (2005) at $\sim 0.3 M_{\odot}$ ($\log M = -0.5$), except for those of González-García et al. 2006 Pinfield et al. 1997) which have no data here.

parable age, size and mass: they have ages of 590^{+150}_{-120} Myr (Fossati et al. 2008) and 625 ± 50 Myr (Bouvier et al. 2008), tidal radii of 11.5 ± 0.3 pc (3.5 ± 0.1 deg, Kraus & Hillenbrand 2007) and 10.3 pc (12.5 deg, Bouvier et al. 2008), and masses of $550 \pm 40 M_{\odot}$ (Kraus & Hillenbrand 2007) and about $400 M_{\odot}$ (Bouvier et al. 2008), respectively. Therefore, we can expect that the potential well is the same (at least today). Only the metallicity may be slightly different, assuming the most recent measurement for Praesepe: $[\text{Fe}/\text{H}] = +0.27 \pm 0.10$ for the latest metallicity measurement of Praesepe (Pace et al. 2008) and $[\text{Fe}/\text{H}] = +0.14 \pm 0.05$ for the Hyades (Bouvier et al. 2008), although a metallicity as low as $+0.038 \pm 0.039$ (Friel & Boesgaard 1992) has been reported for Praesepe. It is unclear how this metallicity difference could explain the significantly different mass functions.

It is a priori possible that different binary mass fractions in Praesepe and the Hyades could account for the difference in their observed (i.e. system, rather than star) mass functions. The binary fraction in Praesepe for different mass intervals was obtained by Pinfield et al. (2003): $17^{+6}_{-4}\%$ for $1.0\text{--}0.6 M_{\odot}$, $31^{+7}_{-6}\%$ for $0.6\text{--}0.35 M_{\odot}$, $44 \pm 6\%$ for $0.35\text{--}0.2 M_{\odot}$ and $47^{+13}_{-11}\%$ for $0.11\text{--}0.09 M_{\odot}$. As for the Hyades, Gizis & Reid (1995) observed a binary fraction of $27 \pm 16\%$ for their sample of stars ($\leq 0.4 M_{\odot}$), which

is consistent with another determination of the Hyades binary fraction of $30 \pm 6\%$ from Patience et al. (1998) (for a primary mass of $\sim 0.6\text{--}2.8 M_{\odot}$). From these figures we see no significant difference in the binary fractions of the two clusters (even if primarily because the uncertainties are quite large), so this cannot be used to explain the difference between in their mass functions. Of course, if the typical mass ratio in a binary system is different in the two clusters then this may be able to account for some difference in the mass functions, but there is also no evidence to support (or refute) this.

A distinction between the two clusters could be the spatial distribution of the members. Indeed, Holland et al. (2000) observed that the Praesepe cluster might be composed of two merged clusters with different ages, one main cluster of $630 M_{\odot}$ and a second subcluster of $30 M_{\odot}$. It was even proposed that faint low-mass members of the subcluster could appear as Praesepe brown dwarf candidates (Chappelle et al. 2005). However, Adams et al. (2002a) did not find evidence of a subcluster in Praesepe. Based on the spatial distribution of the main cluster and subcluster from Holland et al. (2000), our survey only overlaps the main cluster. In addition, a collision between two clusters could not explain alone an increase of the MF down

to $0.1 M_{\odot}$, as such a collision would rather remove low-mass member of the clusters.

By comparing the MF of the Hyades with the one of the Pleiades (~ 120 Myr), Bouvier et al. (2008) concluded that dynamical evolution was responsible for the deficiency observed in the very-low mass star and BD regime in the Hyades. However, this deficiency is not seen in Praesepe. One possible implication is that Praesepe has been less affected by dynamical evolution, i.e. evaporation of low mass members which are expected to have higher speeds based on equipartition of energy. On the other hand, if dynamical evolution *has* affected Praesepe in the same way, then it cannot have had the same initial mass function and/or initial conditions as the Hyades. Dynamical interaction between one of these clusters and another object (such as another open cluster in the past) could explain the discrepancies between the two MFs.

6. Conclusions

We have presented the results of a survey to study the mass function of the old open cluster Praesepe. The survey consisted of optical I_c -band photometry and NIR J and K_s -band photometry with a total coverage of 3.1 deg^2 , down to the substellar regime, with a 5σ detection limit corresponding to $0.05 M_{\odot}$ (the detection completeness to this level is $\sim 87\%$).

Our final sample yields 123 photometric cluster member candidates based on a selection assuming a dust-free atmosphere and 27 photometric cluster candidates based on a selection assuming a dusty atmosphere. We estimate the contamination by field M-dwarfs to be 13% or less. Among our cluster candidates, six objects have theoretical masses equal to or less than the stellar/substellar boundary at $0.072 M_{\odot}$.

We observed that the MF of Praesepe is characterized by a rise in the number of objects from $0.6 M_{\odot}$ down to $0.1 M_{\odot}$, followed by a turn-over in the MF at $\sim 0.1 M_{\odot}$. The rise is in agreement with the Praesepe MFs derived in several previous studies (Hambly et al. 1995; Kraus & Hillenbrand 2007; Baker & Jameson 2009) but disagrees with Adams et al. (2002a).

We have compared the mass function of Praesepe with one derived for the Hyades and have observed a significant difference: while the Hyades has a maximum at $0.35 M_{\odot}$, Praesepe has a maximum at a much lower mass, $0.1 M_{\odot}$. Assuming that they have similar ages (as main sequence fitting suggests), we conclude that the clusters either had different *initial* mass functions or that dynamical interaction has modified the evolution of one or both. More specifically, in the latter case, dynamical evaporation does not seem to have influenced the Hyades and Praesepe in the same way. A difference in the binary fraction or mass ratios could also cause a difference in the mass functions, but determinations of these are not yet precise enough to suggest any difference.

Acknowledgements. SB and CBJ acknowledge support from the Deutsche Forschungsgemeinschaft (DFG) grant BA2163 (Emmy-Noether Programme) to CBJ. SB thanks the Calar Alto observatory staff for support and Kester Smith for observations performed in January 2007. We are grateful to the referee, Nigel Hambly, for his constructive comments and suggestions. We acknowledge Klaus Meisenheimer and Marie-Hélène Nicol for useful discussions about galaxy contamination. IRAF is distributed by the National Optical Astronomy Observatories, which are operated by the Association of Universities for Research in Astronomy, Inc., under cooperative agreement with the National Science Foundation. Some data analysis in this article has made use of the freely available R statistical package, <http://www.r-project.org>. This research has made use of the SIMBAD database, operated at CDS, Strasbourg, France. This publication makes use of data products from the Two Micron All Sky Survey, which

is a joint project of the University of Massachusetts and the Infrared Processing and Analysis Center/California Institute of Technology, funded by the National Aeronautics and Space Administration and the National Science Foundation.

References

- Adams, J. D., Stauffer, J. R., Skrutskie, M. F., et al., 2002a, *AJ*, 124, 1570
- Adams, T., Davies, M. B., Jameson, R. F. & Scally, A., 2002b, *MNRAS*, 333, 547
- Allard, F., Hauschildt, P. H., Alexander, D. R., et al., 2001, *ApJ*, 556, 357
- An, D., Terndrup, D. M., Pinsonneault, M. H., et al., 2007, *ApJ*, 655, 233
- Baade, D., Meisenheimer, K., Iwert, O., et al., 1999, *The Messenger* 95, 15
- Bailer-Jones, C. A. L. & Mundt, R., 2001, *A&A*, 367, 218
- Bailer-Jones, C. A., Bizenberger, P. & Storz, C., 2000, *Society of Photo-Optical Instrumentation Engineers (SPIE) Conference Series*, 4008, 1305
- Baker, D. E. A. & Jameson, R. F., 2009, *Brown Dwarfs in Praesepe: A search for low mass members using UKIDSS*, JENAM2009 meeting, poster no. 3-P03
- Barrado y Navascués, D., Stauffer, J. R. & Jayawardhana, R., 2004, *ApJ*, 614, 386
- Baraffe, I., Chabrier, G., Allard, F. & Hauschildt, P. H., 1998, *A&A*, 337, 403
- Baraffe, I., Chabrier, G., Allard, F. & Hauschildt, P. H., 2002, *A&A*, 382, 563
- Baumeister, H., Bizenberger, P., Bailer-Jones, C. A. L., Kovács, Z., Röser, H.-J. and Rohloff, R.-R., 2003, *Society of Photo-Optical Instrumentation Engineers (SPIE) Conference Series*, 4841, 343
- Béjar, V. J. S., Martín, E. L., Zapatero Osorio, M. R., et al., 2001, *ApJ*, 556, 830
- Boesgaard, A. M. & Budge, K. G., 1988, *ApJ*, 332, 410
- Boudreault, S. & Bailer-Jones, C. A. L., 2009, submitted to the *ApJ* (arXiv:0909.0842v1)
- Bouvier, J., Kendall, T. T., Meeus, G., et al., 2008, *A&A*, 481, 661
- Briceño, C., Luhman, K. L., Hartmann, L., Stauffer, J. R. & Kirkpatrick, J. D., 2002, *ApJ*, 580, 317
- Caballero, J. A., Béjar, V. J. S., Rebolo, R., et al., 2007, *A&A*, 470, 903
- Caballero, J. A., Burgasser, A. J. & Klement, R., 2008, *A&A*, 488, 181
- Chabrier, G., Baraffe, I., Allard, F. & Hauschildt, P., 2000, *ApJ*, 542, 464
- Chabrier, G., 2003, *PASP*, 115, 763
- Chabrier, G., Baraffe, I., Allard, F. & Hauschildt, P. H. 2005, preprint (astro-ph/0509798)
- Chappelle, R. J., Pinfield, D. J., Steele, I. A., et al., 2005, *MNRAS*, 361, 1323
- Close, L. M., Lenz, R., Guirado, J. C., et al., 2005, *Nature*, 433, 286
- Colina, L., Bohlin, R. & Castelli, F., 1996, *Instrument Science Report CAL/SCS*, 8, 1
- D'Antona, F. & Mazzitelli, I., 1985, *ApJ*, 296, 502
- de la Fuente Marcos, R. & de la Fuente Marcos, C., 2000, *Ap&SS*, 271, 127
- Fossati, L., Bagnulo, S., Landstreet, J., et al., 2008, *A&A*, 483, 891
- Friel, E. D. & Boesgaard, A. M., 1992, *ApJ*, 387, 170
- Gizis, J. & Reid, I. N., 1995, *AJ*, 110, 1248
- González-García, B. M., Zapatero Osorio, M. R., Béjar, V. J. S., et al., 2006, *A&A*, 460, 799
- Hambly, N. C., Steele, I. A., Hawkins, M. R. S. & Jameson, R. F., 1995, *MNRAS*, 273, 505
- Hambly, N. C., Hodgkin, S. T., Cossburn, M. R. & Jameson, R. F., 1999, *MNRAS*, 303, 835
- Harris, H. C., Vrba, F. J., Dahn, C. C., et al., 1999, *AJ*, 117, 339
- Hauschildt, P. H., Allard, F. & Baron, E., 1999a, *ApJ*, 512, 377
- Hauschildt, P. H., Allard, F., Ferguson, J., Baron, E. & Alexander, D. R., 1999b, *ApJ*, 525, 871
- Henry, T. J., Jao, W.-C., Subasavage, J. P., et al., 2006, *AJ*, 132, 2360
- Hester, J. J., Scowen, P. A., Sankrit, R., et al., 1996, *AJ*, 111, 2349
- Hillenbrand, L. A. & Carpenter, J. M., 2000, *ApJ*, 540, 236
- Hodgkin, S. T., Pinfield, D. J., Jameson, R. F., et al., 1999, *MNRAS*, 310, 87
- Holland, K., Jameson, R. F., Hodgkin, S., et al., 2000, *MNRAS*, 319, 956
- Howell, S. B. 1989, *PASP*, 101, 616
- Jameson, R. F., Dobbie, P. D., Hodgkin, S. T. & Pinfield, D. J., 2002, *MNRAS*, 335, 853
- Johnstone, D., Hollenbach, D. & Bally, J., 1998, *ApJ*, 499, 758
- Jordi, K., Grebel, E. K. & Ammon, K., 2006, *A&A*, 460, 339
- Kraus, A. L. & Hillenbrand, L. A., 2007, *AJ*, 134, 2340
- Kroupa, P. & Bouvier, J., 2003, *MNRAS*, 346, 369
- Kulkarni, S. & Golimowski, D., 1995, *News Release Number: STSci-1995-48*
- Kumar, M. S. N. & Schmeja, S., 2007, *A&A*, 471, 33
- van Leeuwen, F. 2009, *A&A*, 497, 209
- Lodieu, N., Dobbie, P. D., Deacon, N. R., et al., 2007, *MNRAS*, 380, 712
- Luhman, K. L. & Rieke, G. H., 1999, *ApJ*, 525, 440
- Luhman, K. L., 2000a, *ApJ*, 544, 1044
- Luhman, K. L., 2004, *ApJ*, 617, 1216
- Mac Low, M.-M. & Klessen, R. S., 2004, *Rev. Mod. Phys.*, 76, 125
- Magazzù, A., Rebolo, R., Zapatero Osorio, M. R., et al., 1998, *ApJ*, 497, L77

- Meisenheimer, K., Wolf, C. & Nicol, M.-H., 2009, *in prep.*
- Morau, E., Bouvier, J., Stauffer, J. R. & Cuillandre, J.-C., 2003, A&A, 400, 891
- Muench, A. A., Lada, E. A., Lada, C. J., et al., 2003, AJ, 125, 2029
- Nicol, M.-H., *PhD thesis*, University of Heidelberg (2009)
- Pace, G., Pasquini, L. & Franois, P. 2008, A&A, 489, 403
- Patience, J., Ghez, A. M., Reid, I. N., et al., 1998, AJ, 115, 1972
- Pickett, Brian K., Durisen, Richard H., Cassen, Patrick & Mejia, Annie C., 2000, ApJ, 540, L95
- Pinfield, D. J., Hodgkin, S. T., Jameson, R. F., et al., 1997, MNRAS, 287, 180
- Pinfield, D. J., Dobbie, P. D., Jameson, R. F., Steele, I. A., Jones, H. R. A. & Katsiyannis, A. C. 2003, MNRAS, 342, 1241
- Reid, I. N. & Hawley, S. L., 1999, AJ, 117, 343
- Reipurth, B. & Clarke, C., 2001, AJ, 122, 432
- Salpeter, E. E., 1955, ApJ, 121, 161
- Slesnick, C. L., Hillenbrand, L. A. & Carpenter, J. M., 2004, ApJ, 610, 1045
- Shu, F. H., Adams, F. C. & Lizano, S., 1987, ARA&A, 25, 23
- Stern, D., Kirkpatrick, J. D., Allen, L. E., et al., 2007, ApJ, 663, 677
- Taylor, B. J. 2006, AJ, 132, 2453